

You are cordially invited to attend this educational program



Thrombotic Risk and Symptom Burden in Patients With Polycythemia Vera | Identify When to Intervene With Jakafi®

REGISTRATION



Online <https://sphase.info/inc11534>

You may also register by contacting your Incyte representative(s) Carole Yates at (415) 518-3889 or cyates@incyte.com with the following information: name, title/degree, state(s) and state license number(s), affiliation, address, phone, and email.

Prior to registering, please review the program title and speaker to ensure you have not attended this program before.

Please see Indications, Important Safety Information for Jakafi on back cover and accompanying Full Prescribing Information.

Please note this program is intended for US healthcare professionals who practice in a specialty relevant to the program's FDA-approved indication or disease state. This program is sponsored by Incyte Corporation and is not eligible for CE credits.

This is an educational event intended only for appropriate healthcare professionals. Spouses, guests, and other individuals who are not the intended audience of this educational program are not permitted to attend. Healthcare professionals who are subject to federal, state, or local laws or government ethics restrictions may not attend this event. Incyte will report the cost of any meals provided at this event as required by federal, state, or local laws.

Incyte and its representatives will process your personal information that you provide when you register in order to attend an educational event presented by Incyte. You can learn more about Incyte's privacy practices at the following site: <https://www.incyte.com/privacy-policy>. Please contact privacy@incyte.com if you have any questions or concerns.



PRESENTED BY

Rhonda Hewitt, MSN, ANP, AOCNP

Stanford Cancer Center
Palo Alto, CA



Monday, July 15, 2024

6:30 PM

Pacific Standard Time



Flemings's Prime Steakhouse & Wine Bar

180 El Camino Real
Palo Alto, CA 94304

The program will begin at 6:30 PM. Please plan to arrive 15 minutes early.

Due to a change in policy, Incyte will no longer provide or pay for alcohol at Speaker Programs.

Appropriate attendees include licensed healthcare professionals (HCPs) with a direct role in patient care.

INDICATIONS & USAGE

- Jakafi® (ruxolitinib) is indicated for treatment of polycythemia vera (PV) in adults who have had an inadequate response to or are intolerant of hydroxyurea.
- Jakafi is indicated for treatment of intermediate or high-risk myelofibrosis (MF), including primary MF, post-polycythemia vera MF and post-essential thrombocythemia MF in adults.

WARNINGS & PRECAUTIONS

- Treatment with Jakafi® (ruxolitinib) can cause thrombocytopenia, anemia and neutropenia, which are each dose-related effects. Perform a pre-treatment complete blood count (CBC) and monitor CBCs every 2 to 4 weeks until doses are stabilized, and then as clinically indicated
- Manage thrombocytopenia by reducing the dose or temporarily interrupting Jakafi. Platelet transfusions may be necessary
- Patients developing anemia may require blood transfusions and/or dose modifications of Jakafi
- Severe neutropenia (ANC $<0.5 \times 10^9/L$) was generally reversible by withholding Jakafi until recovery
- Serious bacterial, mycobacterial, fungal and viral infections have occurred. Delay starting Jakafi until active serious infections have resolved. Observe patients receiving Jakafi for signs and symptoms of infection and manage promptly. Use active surveillance and prophylactic antibiotics according to clinical guidelines
- Tuberculosis (TB) infection has been reported. Observe patients taking Jakafi for signs and symptoms of active TB and manage promptly. Prior to initiating Jakafi, evaluate patients for TB risk factors and test those at higher risk for latent infection. Consult a physician with expertise in the treatment of TB before starting Jakafi in patients with evidence of active or latent TB. Continuation of Jakafi during treatment of active TB should be based on the overall risk-benefit determination
- Progressive multifocal leukoencephalopathy (PML) has occurred with Jakafi treatment. If PML is suspected, stop Jakafi and evaluate
- Herpes zoster infection has been reported in patients receiving Jakafi. Advise patients about early signs and symptoms of herpes zoster and to seek early treatment. Herpes simplex virus reactivation and/or dissemination has been reported in patients receiving Jakafi. Monitor patients for the development of herpes simplex infections. If a patient develops evidence of dissemination of herpes simplex, consider interrupting treatment with Jakafi; patients should be promptly treated and monitored according to clinical guidelines
- Increases in hepatitis B viral load with or without associated elevations in alanine aminotransferase and aspartate aminotransferase have been reported in patients with chronic hepatitis B virus (HBV) infections. Monitor and treat patients with chronic HBV infection according to clinical guidelines
- When discontinuing Jakafi, myeloproliferative neoplasm-related symptoms may return within one week. After discontinuation, some patients with myelofibrosis have experienced fever, respiratory distress, hypotension, DIC, or multi-organ failure. If any of these occur after discontinuation or while tapering Jakafi, evaluate and treat any intercurrent illness and consider restarting or increasing the dose of Jakafi. Instruct patients not to interrupt or discontinue Jakafi without consulting their physician. When discontinuing or interrupting Jakafi for reasons other than thrombocytopenia or neutropenia, consider gradual tapering rather than abrupt discontinuation
- Non-melanoma skin cancers (NMSC) including basal cell, squamous cell, and Merkel cell carcinoma have occurred. Perform periodic skin examinations
- Treatment with Jakafi has been associated with increases in total cholesterol, low-density lipoprotein cholesterol, and triglycerides. Assess lipid parameters 8-12 weeks after initiating Jakafi. Monitor and treat according to clinical guidelines for the management of hyperlipidemia
- Another JAK-inhibitor has increased the risk of major adverse cardiovascular events (MACE), including cardiovascular death, myocardial infarction, and stroke (compared to those treated with tumor TNF blockers) in patients with rheumatoid arthritis, a condition for which Jakafi is not indicated. Consider the benefits and risks for the individual patient prior to initiating or continuing therapy with Jakafi particularly in patients who are current or past smokers and patients with other cardiovascular risk factors. Patients should be informed about the symptoms of serious cardiovascular events and the steps to take if they occur
- Another JAK-inhibitor has increased the risk of thrombosis, including deep venous thrombosis (DVT), pulmonary embolism (PE), and arterial thrombosis (compared to those treated with TNF blockers) in patients with rheumatoid arthritis, a condition for which Jakafi is not indicated. In patients with myelofibrosis (MF) and polycythemia vera (PV) treated with Jakafi in clinical trials, the rates of thromboembolic events were similar in Jakafi and control treated patients. Patients with symptoms of thrombosis should be promptly evaluated and treated appropriately
- Another JAK-inhibitor has increased the risk of lymphoma and other malignancies excluding NMSC (compared to those treated with TNF blockers) in patients with rheumatoid arthritis, a condition for which Jakafi is not indicated. Patients who are current or past smokers are at additional increased risk. Consider the benefits and risks for the individual patient prior to initiating or continuing therapy with Jakafi, particularly in patients with a known secondary malignancy (other than a successfully treated NMSC), patients who develop a malignancy, and patients who are current or past smokers
- In myelofibrosis and polycythemia vera, the most common nonhematologic adverse reactions (incidence $\geq 15\%$) were bruising, dizziness, headache, and diarrhea. In acute graft-versus-host disease, the most common nonhematologic adverse reactions (incidence $>50\%$) were infections (pathogen not specified) and edema. In chronic graft-versus-host disease, the most common nonhematologic adverse reactions (incidence $>20\%$) were infections (pathogen not specified) and viral infections
- Avoid concomitant use with fluconazole doses greater than 200 mg. Dose modifications may be required when administering Jakafi with fluconazole doses of 200 mg or less, or with strong CYP3A4 inhibitors, or in patients with renal or hepatic impairment. Patients should be closely monitored and the dose titrated based on safety and efficacy
- Use of Jakafi during pregnancy is not recommended and should only be used if the potential benefit justifies the potential risk to the fetus. Women taking Jakafi should not breastfeed during treatment and for 2 weeks after the final dose

Please see accompanying Full Prescribing Information for Jakafi.